

Emotion Recognition in Objects in Patients With Neurological Disease

Michelle N. Shiota
Arizona State University

Michaela L. Simpson
University of California, Berkeley

Heidi E. Kirsch
University of California, San Francisco

Robert W. Levenson
University of California, Berkeley



Objective: Considerable research indicates that individuals with dementia have deficits in the ability to recognize emotion in other people. The present study examined ability to detect emotional qualities of objects. **Method:** Fifty-two patients with frontotemporal dementia (FTD), 20 patients with Alzheimer’s disease (AD), 18 patients awaiting surgery for intractable epilepsy, and 159 healthy controls completed a newly developed test of ability to recognize emotional qualities of art (music and paintings), and pleasantness in simple sensory stimuli (tactile, olfactory, auditory), and to make aesthetic judgments (geometric shapes, room décor). A subset of participants also completed a test of ability to recognize emotions in other people. **Results:** Patients with FTD showed a marked deficit in ability to recognize the emotions conveyed in art, compared with both healthy individuals and patients with AD (relative to controls, deficits in patients with AD only approached significance). This deficit remained robust after controlling for FTD patients’ ability to recognize pleasantness in simple sensory stimuli, make aesthetic judgments, identify odors, and identify emotions in other people. Neither FTD nor AD patients showed deficits in recognizing pleasant sensory stimuli or making aesthetic judgments. Exploratory analysis of patients with epilepsy revealed no deficits in any of these domains. **Conclusion:** Patients with FTD (but not AD) showed a significant, specific deficit in ability to interpret emotional messages in art, echoing FTD-related deficits in recognizing emotions in other people. This finding adds to our understanding of the impact these diseases have on the lives of patients and their caregivers.

General Scientific Summary

The ability to recognize emotions in others, and to enjoy pleasing sensations and aesthetic environments, is an important part of human life. Using a newly developed test of the ability to recognize emotional qualities in art, pleasantness in sensory stimuli, and to make aesthetic judgments, we found deficits in these abilities in patients with two kinds of dementia (most profoundly in frontotemporal dementia and less so in Alzheimer’s disease) but not in patients with epilepsy. With the aging population and increasing number of families caring for loved ones with dementia, it is important to understand how these diseases affect this important aspect of emotional functioning.

Keywords: emotion recognition, aesthetic judgments, frontotemporal dementia, Alzheimer’s disease, epilepsy

Supplemental materials: <http://dx.doi.org/10.1037/neu0000587.supp>

This article was published Online First September 2, 2019.

Michelle N. Shiota, Department of Psychology, Arizona State University; Michaela L. Simpson, Department of Psychology, University of California, Berkeley; Heidi E. Kirsch, Department of Neurology, University of California, San Francisco;  Robert W. Levenson, Department of Psychology, University of California, Berkeley.

This research was supported by National Institute of Health Grants R01 AG041762, AG019724, and K23 NS047100.

 The experiment materials are available at https://osf.io/7b4sc/?view_only=6872f94c39a749a8a701e971215b5d16.

Correspondence concerning this article should be addressed to Robert W. Levenson, Department of Psychology, University of California, 3206 Berkeley Way West, Berkeley, CA 94720-1650. E-mail: boblev@berkeley.edu

Recognizing and responding to the emotional properties of environmental stimuli is important for successfully navigating the external world and can play a valuable role in supporting well-being. This broad category of functions includes several related but distinguishable abilities. At the simplest level, people can enjoy pleasant simple stimuli, such as the soft texture of a cashmere blanket or the scent of a rose. People also respond to more complex aesthetic features of the environment, such as the harmony of colors in a room. At a more sophisticated level, people can recognize emotions expressed by others through facial expressions and tone of voice, and can even detect emotional messages communicated through the formal features of art (e.g., perceiving a piece of music as “happy” or “sad”). These abilities can provide much enjoyment in life, and even contribute to resilience and well-being during times of stress (e.g., Folkman, 1997; Shiota, 2006).

Much research has documented deficits among individuals with frontotemporal dementia (FTD) in aspects of empathy toward other people, including the ability to recognize their emotions (e.g., Goodkind et al., 2015; Rosen et al., 2004). Less is known about the implications of FTD and other neurological disorders for responding to emotionally relevant properties of objects in the environment such as art, aesthetic judgments, and simple sensory stimuli. The current study aims to help address this gap.

Emotional Responses to Aesthetics and Art

Human responses to the emotional properties of objects in the environment can take two broad forms: appreciating the object’s sensory or aesthetic pleasantness (vs. unpleasantness); and decoding its emotional message or tone (i.e., the emotion it communicates). While people can evaluate even very simple stimuli in terms of pleasantness, such as geometric shapes and scents, works of art (including music and painting) are often intended by the artist to convey a specific emotional quality, such as joy or anguish. Abilities to evaluate stimulus pleasantness and to decode emotional content in art overlap to some extent, but reflect distinct adaptive functions, and may be supported by distinct psychological and neural mechanisms.

The sensory stimuli we identify as pleasing are typically those that helped sustain the lives of our ancestors, activating approach behaviors that are critical for adaptive fitness (Shiota et al., 2017). For example, the human sense of smell is closely linked to the sense of taste (Shepherd, 2006), and the most strongly preferred scents cross-culturally are those associated with palatable food (Gang, 2005). Similarly, people prefer soft, fuzzy textures rather than rough ones (Essick et al., 2010), reflecting an innate preference for the feel of mammalian parents’ skin and fur (Harlow, 1958). In the domain of sound, people prefer individual tones and consonant (vs. dissonant) combinations of tones that are characteristic of human speech (Schwartz, Howe, & Purves, 2003). In the visual domain, people tend to prefer figures that are symmetrical (an indicator of health and developmental stability in living organisms; Møller, 1997); moderately complex, with sharper angles between sides and a combination of concave and convex angles (Friedenberg & Bertamini, 2015); and perceptually balanced around the geometric center of the image (Friedenberg & Bertamini, 2015).

In contrast, the ability to perceive emotional messages in art may be a byproduct of our ability to decode emotional expressions in other people and use metaphors to describe emotional feelings. For example, the acoustic properties of music that convey emotion have been found to correspond to those of the human voice that serve the same purpose (Juslin & Laukka, 2003). In the visual arts, simple properties of color and line are perceived as expressing affect in the same two-dimensional valence and arousal space people use to describe their own feelings (Collier, 1996). Even in verbal language, metaphors involving simple sensory experiences are more effective at evoking emotions than literal statements with the same meaning (Citron & Goldberg, 2014). The ability to associate simple stimulus properties with emotional meaning through metaphor, combined with the ability to perceive the emotion expressed by a being other than the self, may combine to produce the ability to decode the emotional meaning of art.

The available research on neural structures associated with these functions suggests some overlap, and some differentiation. A rich body of research identifies opioid receptor-rich regions of the nucleus accumbens and ventral pallidum as highly responsive to sensory enjoyment or “liking” of simple stimuli such as pleasant tastes (Berridge & Kringelbach, 2013). Studies of neural activity during judgments of visual aesthetic preference, whether for simple geometric shapes or complex paintings, identify the orbitofrontal cortex (OFC) as an area that is more active while viewing preferred stimuli (e.g., Jacobsen, Schubotz, Höfel, & Cramon, 2006; Kawabata & Zeki, 2004). This is consistent with the evidence implicating OFC activation in stimulus valuation more generally, such as in the context of economic games, assessments of physical beauty, and aesthetic judgments (Chatterjee, 2011; FitzGerald, Seymour, & Dolan, 2009; Stalnaker, Cooch, & Schoenbaum, 2015). Multiple studies have also documented greater activity in anterior cingulate cortex (ACC) while viewing preferred relative to nonpreferred paintings (Kawabata & Zeki, 2004; Vartanian & Goel, 2004).

Meta-analysis results suggest that processing of linguistic metaphor, relative to literal language, is associated with heightened activation in the inferior frontal gyrus (IFG), the amygdala, and the superior temporal gyrus (STG), among other regions (Bohrn, Altmann, & Jacobs, 2012; Citron & Goldberg, 2014). The latter of these are noteworthy as heightened amygdala and STG activation are also commonly seen in emotion recognition tasks with human face stimuli (Adolphs, 2002). Empirical reviews and meta-analysis results further suggest that the amygdala, anterior insula, OFC, and regions of the ACC and medial prefrontal cortex show heightened activation during empathy tasks, although the exact profile of activation depends on specific task demands (e.g., infer vs. feel the target’s emotion; Adolphs, 2002; Fan, Duncan, de Greck, & Northoff, 2011; Zaki & Ochsner, 2012).

Emotion Processing in Neurological Patients

Neurodegenerative diseases can affect brain circuitry that is critical for emotion processing (Rosen & Levenson, 2009). A number of studies have found that individuals with Alzheimer’s disease (AD) and frontotemporal dementia (FTD) have deficits in the ability to recognize emotion in other people (e.g., Fernandez-Duque & Black, 2005; Goodkind et al., 2015; Hargrave, Maddock, & Stone, 2002; Hsieh, Hodges, & Piguet, 2013; Keane, Calder,

Hodges, & Young, 2002; Lavenu, Pasquier, Lebert, Petit, & Van der Linden, 1999; Ostos, Schenk, Baenziger, & von Gunten, 2011; Rosen et al., 2004; Rosen, Perry et al., 2002; Weiss et al., 2008; Werner et al., 2007). When patients with AD and FTD have been compared, these deficits appear to be more pronounced in FTD (Goodkind et al., 2015; Lavenu et al., 1999). Among other neurological diseases, deficits in ability to recognize emotion in people have also been reported in individuals with epilepsy (Meletti et al., 2003; Reynders, Broks, Dickson, Lee, & Turpin, 2005) and amygdala lesions (Adolphs et al., 1999). In addition, similar deficits have been found in individuals with a number of psychiatric disorders (e.g., Ashwin, Chapman, Colle, & Baron-Cohen, 2006; Chan, Li, Cheung, & Gong, 2010; Rocca, van den Heuvel, Caetano, & Lafer, 2009).

One of the intriguing findings in this literature has been that individuals with dementia have deficits in recognizing negative emotions in other people, but that the ability to recognize positive emotions may be preserved (Fernandez-Duque & Black, 2005; Kessels et al., 2007; Lavenu et al., 1999; Lough et al., 2006). Although evolutionary accounts of emotion often emphasize the importance of negative emotions (which help us survive when confronted with threats), the ability to detect positive emotions is extremely important as well. Expressions of positive emotion signal potential relationship partners who may be committed to our welfare and able to provide important resources (e.g., Gonzaga, Keltner, Londahl, & Smith, 2001). Given the needs of individuals with dementia for care and assistance by others, preservation of the ability to recognize positive emotion would be important to document. However, as we have suggested elsewhere (Goodkind et al., 2015), findings of preserved recognition of positive emotions may result from a structural asymmetry in most tests of emotion recognition. These tests typically utilize photographs of emotional facial expressions that have multiple negative emotions but only a single positive emotion (happiness portrayed by a smiling face), thus making the detection of positive emotions much easier. Utilizing a film-based emotion identification test that assessed multiple positive (e.g., affection, amusement), negative (e.g., anger, fear), and self-conscious (e.g., embarrassment, pride) emotions, we (Goodkind et al., 2015) found deficits in recognition for all three kinds of emotion in patients with FTD.

A much smaller group of studies has examined dementia patients' ability to identify the emotional qualities of nonface targets, such as simple sensory stimuli and works of art (e.g., Boutoleau-Bretonnière et al., 2016; Cohen et al., 2016; Dellacherie, Ehrlé, & Samson, 2008; Omar et al., 2011). For example, Boutoleau-Bretonnière and colleagues (2016) found that FTD patients generally rated abstract paintings as less pleasant and less emotionally moving than did nonpatient controls. Cohen and colleagues (2016) found that FTLT patients received lower scores than controls on a match-to-sample task relying on the affective valence of abstract art stimuli, with normative valence established through pilot research. Although these studies consistently identify deficits in FTD patients relative to controls, it is not clear from the tasks whether the deficit lies in evaluating the pleasantness/enjoyability of stimuli (i.e., which is more pleasant?), in decoding the emotional message of the stimuli (i.e., which communicates happiness?), or both. The present study uses a novel test that helps tease these two processes apart.

Comparing Emotion Recognition in Different Neurological Disorders

As noted above, most studies of the ability of dementia patients to recognize emotion in people have been conducted with FTD or AD patients. FTD is a neurodegenerative disease that causes atrophy in the frontal and temporal lobes (Zhang et al., 2009). In the initial stages of the disease, FTD patients show profound changes in emotion, personality, and behavior, including loss of empathy, loss of social awareness, disinhibition, apathy, loss of insight, and impaired emotion recognition (Rankin, Kramer, & Miller, 2005; Snowden et al., 2001). In contrast, AD causes atrophy in the medial temporal and parietal lobes (Zhang et al., 2009). In the initial stages, AD patients show marked deterioration of memory and cognitive abilities, with relative preservation of emotional functioning (e.g., Bucks & Radford, 2004; Goodkind, Gyurak, McCarthy, Miller, & Levenson, 2010; Goodkind et al., 2015; Lavenu et al., 1999). Although epilepsy is not considered to be a form of dementia, it is often accompanied by emotional changes (e.g., depression and anxiety; Hixson & Kirsch, 2009; Kanner, 2009). Moreover, seizure foci in epilepsy can occur in regions (e.g., temporal lobes) that are vulnerable in frontotemporal dementia (Zhang et al., 2009). Despite the obvious virtue of comparing multiple neurological conditions, we are not aware of previous studies of emotion recognition that included FTD, AD, and epilepsy patients as well as healthy controls.

The Present Study

The present study assessed the ability of neurological patients with FTD and AD to recognize emotion in art (paintings and music) and pleasantness in simple sensory stimuli (olfactory, auditory, tactile), and to make aesthetic judgments (geometric shapes, room décor), in comparison to a group of healthy controls, and, in an exploratory analysis, a group of epilepsy patients awaiting surgery for intractable seizures. In a subgroup of the patients with FTD and AD, the ability to recognize emotion in facial expressions was also assessed.

Method

Participants

Participants were 52 patients with FTD (25 males, 27 females), 20 patients with AD (11 males, 9 females), and 159 controls (82 males, 76 females, 1 who declined to indicate gender). Among the FTD patients, 25 were diagnosed with behavioral variant FTD, 21 with semantic dementia, and six with primary nonfluent aphasia. Patients with FTD and AD were recruited through the Memory and Aging Center of the University of California, San Francisco (UCSF), where they received comprehensive neurological and neuropsychological testing. Because significant numbers of participants were assessed prior to the appearance of newer diagnostic criteria (e.g., Gorno-Tempini et al., 2011; Rascovsky et al., 2007, 2011), the Neary clinical criteria (Neary et al., 1998) were used to diagnose FTD. Similarly, the National Institute of Neurological and Communication Disorders criteria and those of the Stroke/Alzheimer's disease and Related Disorders Association (McKhann et al., 1984) were used to diagnose AD. Controls were adult caregivers (mostly spouses) of patients with FTD and AD who were participating in a study of emotional functioning in dementia

caregivers conducted at the Berkeley Psychophysiology Laboratory at the University of California, Berkeley.

An additional group of 18 patients with intractable epilepsy (6 males, 12 females) was recruited by one of the authors (HEK) at the UCSF Medical Center. These patients were all awaiting surgery to alleviate medically refractory seizures. Including patients with intractable focal epilepsy provides an important test of the specificity of findings by establishing whether deficits found in patients with dementia generalize to another neurological condition.

We included patients with AD, different FTD subtypes, and epilepsy to increase the anatomical and behavioral variability in our sample. AD is characterized by neurodegeneration in hippocampus, entorhinal cortex, and posterior cingulate and may leave emotional functioning relatively intact in the early stages of disease (Goodkind et al., 2010; Lavenu & Pasquier, 2005). Compared to AD, FTD is characterized by degeneration in more anterior brain regions and produces more profound deficits in emotional functioning (Levenson & Miller, 2007). Among the three subtypes of FTD included in our sample: (a) behavioral variant FTD is characterized by bilateral frontal lobe degeneration and significant changes in emotion and personality; (b) semantic dementia is characterized by atrophy in anterior temporal areas and language problems; and (c) primary nonfluent aphasia is characterized by predominantly left-sided frontotemporal atrophy and problems with speaking (Neary et al., 1998; Rosen, Gorno-Tempini, et al., 2002; Rosen, Kramer, et al., 2002). Patients with epilepsy who receive surgical treatment for intractable seizures often have seizure foci in the temporal region and may experience significant emotional changes (Meletti et al., 2003; Reynders et al., 2005). Network dysfunction in these patients with epilepsy is thought to involve frontotemporal regions, anatomic substrates that are similar to those affected in FTD. In our sample of 18 epilepsy patients, 15 had established temporal foci with nine having a primary left hemisphere focus and nine having a primary right hemisphere focus.

Patients with dementia and epilepsy and controls in this sample participate in various research projects conducted at UCSF and at UC Berkeley. AD and FTD patients and controls in this sample all participated in a comprehensive assessment of emotional functioning (Levenson, 2007) conducted at UC Berkeley. Some data from these assessments have been published previously (e.g., Goodkind et al., 2015). However, no data from these participants related to detecting the emotional qualities of objects have been published previously.

Measures

Emotion in objects. We utilized the Test of Emotional Aesthetics (TEA), a 16-item test developed at the Berkeley Psychophysiology Laboratory, to assess ability to recognize the emotional qualities of art (paintings, music), simple sensory stimuli (olfactory, auditory, tactile), and to make aesthetic judgments (room décor, geometric shapes). The instructions and visual and auditory stimuli for the TEA are presented on a laptop computer.

Stimuli for the art items included two pairs of paintings and two pairs of short musical excerpts; stimuli within each pair were always by the same artist. In the paintings (Picasso: “Weeping Woman” and “The Dream”; Van Gogh: “Wheat Field with Cy-

presses” and “Wheat Fields with Crows”), each painting in the pair depicts a similar subject through very different formal features, thereby conveying distinct emotional messages. Specifically, whereas “The Dream” and “Wheat Field with Cypressess” use lighter, primary, saturated colors and smoother, curved lines to convey calm positive affect, “Weeping Woman” and “Wheat Fields with Crows” both use darker, nonprimary, less saturated colors and sharper, more jagged lines to convey negative affect (Collier, 1996). Similarly, pairs of musical excerpts (*Living Out Loud* soundtrack, “Ecstasy” and “She’s 34”; Bach Goldberg Variations: #21 and #22) present the same musical theme, one with a quick tempo and major key to convey happiness, and the other with slower tempo and minor key to convey sadness (Gabrielsson & Juslin, 2003). In each art item, the participant was asked to identify which of the pair was “more happy,” relying on the ability to decode the emotional messages of the two options.

Eleven items assessed preferences for simple sensory or aesthetic stimuli: (a) Olfactory 1—scratch and sniff: apple versus natural gas; (b) Olfactory 2—scratch and sniff: mint versus paint thinner; (c) Auditory 1 and 2—consonant versus dissonant piano chords; (d) Tactile 1—reach into bags: Brillo pad versus silk; (e) Tactile 2—reach into bags: cotton versus sandpaper; (f) Aesthetic visual judgments 1 and 2—two rooms decorated with different color schemes; and (j) Aesthetic visual judgments 3, 4, and 5—pairs of geometric and abstract shapes, differing on complexity and balance (one of these was taken from the Visual Aesthetic Sensitivity Test; Götz, Boris, Lynn, & Eysenck, 1979). In each pair of items, the correct response was based on the empirically supported principles of aesthetic preference discussed in the introduction section of this paper. Figure 1 presents example stimulus pairs from the aesthetic visual items. In each preference item, participants are asked which of the two options is “more pleasing.” The 16th item assesses Odor Identification, presenting a single scratch and sniff rose smell with three label choices: smoke, rose, or lemon.

Each of the 16 items was scored correct (1) or incorrect (0), and normative responses were validated with an independent group of healthy controls. Percentage correct scores were computed for the following subscales: (a) Art (four items); Sensory stimuli (six items); Aesthetic visual judgments (five items); and Odor identification (one item). Copies of the TEA (along with instructions for creating the tactile and odor items) are available from the authors.

Emotion in people. The Empathy Film Test (EFT; Goodkind et al., 2015) assesses ability to recognize emotions in other people. It consists of 11 excerpts (approximately 40 s in length) from commercial films each featuring a character expressing one of four positive (affection, amusement, calm, enthusiasm), four negative (anger, disgust, fear, sadness), or three self-conscious (embarrassment, shame, pride) emotions. After each clip, the participant is shown a photo of the main character with a neutral facial expression and a list of the 11 target emotions and asked: “What emotion did the character feel most strongly?” Reliabilities of the EFT subscales (Cronbach’s alpha: positive emotions = .63; negative emotions = .72, self-conscious emotions = .65) have been published previously (Goodkind et al., 2015).

The EFT has several advantages over traditional tests of emotion recognition (e.g., Bowers, Blonder, & Heilman, 1991) in which respondents have to provide the correct emotion label (e.g., “happy”) for a static photograph of an isolated emotional facial




Figure 1. Sample visual aesthetics item stimulus pairs from test of emotional aesthetics. A. Rooms; B. Geometric shapes. In each item, the participant is asked to identify which of the pair is “more pleasing.” Correct answers are (A) left; (B) right. See the online article for the color version of this figure.

expression (e.g., a single smiling person). The EFT has arguably greater ecological validity by virtue of assessing respondents’ ability to recognize emotion using dynamic, multimodal (visual, auditory), and interpersonal stimuli. These conditions more closely mirror real-world emotional judgments compared to photographs, which are static, convey only visual information, and are usually not depicted in interpersonal situations (Goodkind et al., 2015). Moreover, photograph-based tests typically assess multiple negative emotions (e.g., anger, fear, sadness, disgust) but only assess a single positive emotion (happiness as portrayed by a smiling face) and do not assess self-conscious emotions. In contrast, the EFT assesses multiple positive and multiple self-conscious emotions in addition to multiple negative emotions. Thus, in addition to its ecological validity, the EFT can provide a more comprehensive test of the ability to recognize emotion in people.

Each response on the EFT was scored correct (1) or incorrect (0) and a percent correct score was determined for the following subscales: (a) Positive emotion (four items), (b) Negative emotion (four items), and (c) Self-conscious emotion (three items). Because the assessment of ability to recognize emotion in the present study focused primarily on positive emotion (i.e., happiness in art, pleasantness in sensory stimuli and aesthetic judgments), we used the Positive emotion score from the EFT in our primary analyses to control for deficits in emotion recognition in people when assessing deficits in emotion recognition in objects using the TEA. In secondary analyses, we also used the Negative and Self-conscious emotion scores from the EFT.

Procedure

Patients with FTD and AD participated in a comprehensive assessment of emotional functioning (Levenson, 2007) at the Berkeley Psychophysiology Laboratory. As part of this assessment, they completed the TEA and the EFT. Caregivers who accompanied the patients also completed the TEA and the EFT. Epilepsy patients completed the TEA at the UCSF Medical Center but did not complete the EFT. All participants gave informed consent prior to participation. The study was approved by the Committee for the Protection of Human Subjects at the University of California, Berkeley and the University of California, San Francisco.

Data Analysis

Analyses initially focused on the FTD and AD patients and controls because these groups had completed both the TEA and the EFT. An overall 3×4 (Diagnosis [FTD, AD, controls] \times TEA subscale [Art, Sensory stimuli, Aesthetic judgments, Odor identification]) repeated measures MANOVA was computed using the General Linear Model procedure in SPSS Version 24. In this MANOVA, TEA subscale was treated as a repeated measure and all main effects and interactions were analyzed using multivariate *F* tests (which obviate the need to control for violations of sphericity in the repeated measures). Significant effects were followed up by comparisons among means using least-significance difference tests and 95% confidence intervals (CI).

Results

Preliminary Analyses

Psychometrics. Because the TEA has not been used in prior published research, we conducted basic psychometric analyses using our sample of 159 healthy controls. These revealed that the internal consistency of the 16-item test was moderate (Cronbach’s $\alpha = .45$). We expect this reflects both restriction in range in our neurologically healthy sample and the TEA’s assessment of different kinds of emotional judgments. Consistent with range restriction, for controls the percent correct score averaged across all 16 items was 84% ($SE = .89$, range = 44–100%) with the lowest percent correct generally found for the aesthetic judgments of geometric shapes. An item-level reliability analysis revealed that the overall reliability of the scale was not markedly changed by removing any of the individual items.

Sample characteristics. The average age for the sample of patients with FTD and AD and healthy controls was 62.4 years ($SE = 0.6$). This sample was well-educated (mean years of education = 16.5, $SE = 0.2$) and primarily Caucasian (88.0%). Analyses revealed no gender, $\chi^2(2) = 0.35$, $p = .84$, age, $F(2, 226) = .02$, $p = .99$, or education, $F(2, 224) = 1.43$, $p = .24$ differences among the three groups. The sample of epilepsy patients used in the exploratory analysis was younger (mean age = 35.5, $SE = 2.9$) and less educated (mean years of education = 13.7, $SE = 0.4$).

Using data obtained from neuropsychological testing of patients with FTD and AD conducted at UCSF, Mini-Mental State (Folstein, Folstein, & McHugh, 1975) scores (which can range from 0–30, with lower scores indicating greater cognitive impairment)

indicated moderate cognitive impairment ($M = 23.1$, $SE = 0.7$), and clinical dementia rating (Morris, 1993) sum of boxes scores (which can range from 0 to 18, with higher scores indicating greater dementia severity) indicated moderate levels of dementia severity (mean sum of boxes = 5.0, $SE = 0.4$). Table 1 presents demographic and clinical data for the diagnostic groups.

Emotion in Objects

FTD, AD, and controls. The MANOVA of the TEA revealed significant main effects for Diagnosis, $F(2, 228) = 21.96$, $p < .001$ and TEA Subscale, $F(3, 226) = 16.88$, $p < .001$ as well as a significant interaction of Diagnosis \times TEA Subscale, $F(6, 454) = 7.80$, $p < .001$. Examining the main effect for Diagnosis, both FTD patients and AD patients scored lower than controls (FTD vs. controls: 95% CI = -17.15 , -8.86 , $p < .001$; AD vs. controls: 95% CI = -16.73 , -4.41 , $p = .001^1$). Differences between FTD and AD patients were not significant ($p = .483$). Examining the main effect for TEA Subscale, performance on the Art subscale and Sensory stimuli subscale were higher than the Aesthetic judgments subscale (Art vs. Aesthetic judgments: 95% CI = 8.38, 17.78, $p < .001$; Sensory stimuli vs. Aesthetic judgments: 95% CI = 6.40, 15.27, $p < .001$) and Odor identification subscale (Art vs. Odor identification: 95% CI = 10.63, 24.99, $p < .001$; Sensory stimuli vs. Odor identification: 95% CI = 8.66, 22.47, $p < .001$).

The Diagnosis \times Tea Subscale interaction was decomposed by computing univariate Diagnosis ANOVAs for each TEA subscale. These revealed that the main effect for Diagnosis was not significant for Sensory stimuli, $F(2, 228) = 0.57$, $p = .57$ or for Aesthetic judgments, $F(2, 230) = 0.20$, $p = .82$. However, the main effect for Diagnosis was significant for Art, $F(2, 228) = 23.08$, $p < .001$ and Odor identification, $F(2, 228) = 19.01$, $p < .001$. For Art, FTD patients scored lower than AD patients (95% CI = -18.86 , -2.30 , $p = .013$) and lower than controls (95% CI = -22.29 , -12.23 , $p < .001$). AD patients also scored lower than controls, but this difference only approached significance (95% CI = -14.15 , $.79$, $p = .079$). For Odor identification, both dementia patient groups scored lower than controls (FTD vs. controls: 95% CI = -42.57 , -19.33 , $p < .001$; AD vs. controls: 95% CI = -52.82 , -18.31 , $p < .001$), but did not differ from each other (FTD vs. AD: 95% CI = -14.52 , 23.75 , $p = .635$). Table 2 presents TEA data for the diagnostic groups.

Comparing FTD subtypes. The sample sizes for FTD patients with behavioral variant FTD ($N = 25$) and semantic dementia ($N = 21$) were not large. Nonetheless, we thought it useful to conduct analyses to assess performance of these two FTD subtypes on the TEA and its subscales. Using univariate ANOVAs, the two FTD subtypes did not differ on overall TEA scores or on any of the subscales (Overall: $F(1, 44) = .25$, $p = .62$; Art: $F(1, 44) = .57$, $p = .46$; Sensory stimuli: $F(1, 44) = .40$, $p = .53$; Aesthetic judgments: $F(1, 44) = .12$, $p = .73$; Odor identification: $F(1, 44) = .01$, $p = .94$). Moreover, there was no consistent pattern of one subtype performing better across the subscales. The smallish sample sizes require caution in interpreting this null finding, but the pattern of results does not suggest that deficits on the TEA in the larger FTD sample were driven by deficits in a particular subgroup. Table 3 presents TEA scores for the two FTD subtypes.

Summary. Both FTD and AD patients showed lower performance overall on the TEA compared to controls. In terms of

specific deficits, both FTD and AD patients showed deficits in Odor identification but no deficits in detecting pleasantness in Sensory stimuli or Aesthetic judgments. On the Art subscale, FTD patients showed significant deficits, and AD patients showed marginal deficits, compared to healthy controls. An exploratory analysis comparing TEA performance in two FTD subtypes found no evidence of differences between patients with behavioral variant FTD and semantic dementia.

Robustness of FTD Deficits in Recognizing Emotion in Art

Because not all of the participants who completed the TEA completed the EFT, this analysis was conducted using data from a smaller pool of participants (34 FTD, 11 AD, 62 controls) who completed both tests. In this smaller sample, an ANOVA of performance on the Positive emotion subscale of the EFT revealed a significant main effect for Diagnosis, $F(2, 104) = 16.61$, $p < .001$ with both FTD patients and AD patients scoring lower than healthy controls (FTD vs. controls: 95% CI = -34.15 , -15.03 , $p < .001$; AD vs. controls: 95% CI = -42.70 , -13.38 , $p < .001$). Table 2 presents EFT data for the diagnostic groups.

To evaluate the specificity of the deficits in recognizing emotion in art found for FTD patients, we determined whether this deficit was accounted for by deficits in ability to recognize emotion in other kinds of objects and in people. We conducted an analysis of covariance (ANCOVA) examining diagnostic group differences in performance on the Art subscale of the TEA, while controlling for the three other TEA subscales (Sensory stimuli, Aesthetic judgments, Odor identification) and for the Positive emotion subscale of the EFT. This ANCOVA revealed a main effect for diagnostic group that approached significance, $F(2, 106) = 2.97$, $p = .056$. Follow-up comparisons revealed that FTD patients (estimated mean % = 81.85) still scored significantly lower than controls (estimated mean % = 91.51; 95% CI = -17.98 , -1.34 , $p = .023$), even when controlling for the four covariates.

In exploratory analyses, we determined whether FTD patients' deficit in recognizing emotion in art would remain when controlling for their ability to recognize emotion in other objects and their ability to recognize either negative emotions or self-conscious emotions in people. We computed two ANCOVAs that controlled for the three other TEA subscales (Sensory stimuli, Aesthetic judgments, and Odor identification) as well as performance on either the Negative emotion subscale or the Self-conscious emotion subscale of the EFT. In both analyses, controlling for all four covariates, FTD patients still scored significantly lower than controls (Negative emotion on EFT: 95% CI = -19.73 , -2.435 , $p = .013$; Self-conscious emotion on EFT: 95% CI = -17.279 , $-.520$, $p = .038$).

Are Deficits in Recognizing Emotion in Objects Also Found in Patients With Epilepsy?

To determine whether our TEA findings with FTD and AD patients generalized to another patient group, we recomputed our

¹ When the Odor identification item was removed from this analysis, the pairwise contrast between AD patients and controls was no longer significant (CI = -7.67 , 3.19 , $p = .42$). All other results are unchanged.

Table 1
Demographic and Clinical Information for Diagnostic Groups

Variable	FTD	AD	Epilepsy	Controls	Total
<i>N</i>	52	20	18	159	249
Sex: % female	51.9	45.0	66.7	47.8	50.0
Age: Mean (<i>SE</i>)	62.6 (1.0)	62.2 (2.0)	35.5 (2.9)	62.4 (.8)	60.5 (.7)
Education: Years (<i>SE</i>)	16.0 (.4)	17.0 (.8)	13.7 (.4)	16.5 (.2)	16.3 (.2)
Race/Ethnicity: %					
Caucasian	86.2	88.9	n/a	88.5	88.0
Asian/Asian American	9.8	5.6	n/a	3.2	4.9
Hispanic	3.9	0	n/a	1.9	2.2
African American	0	0	n/a	1.3	.9
Native Hawaiian/Pacific Islander	0	0	n/a	1.3	.9
American Indian/Native Alaskan	0	0	n/a	.6	.4
Refused to state/Unknown	0	5.6	n/a	0	.4
Clinical information: Mean (<i>SE</i>)					
Mini-Mental State Examination	24.1 (.8)	20.9 (1.0)	n/a	n/a	23.1 (.7)
Clinical dementia rating sum of boxes	4.9 (.5)	5.3 (.5)	n/a	n/a	5.0 (.4)

Note. FTD = frontotemporal dementia; AD = Alzheimer's disease; n/a = not available; *SE* = standard error.

overall MANOVA adding data from the 18 patients with epilepsy. In this expanded ANOVA there were significant effects for Diagnosis, $F(3, 245) = 15.97, p < .001$, TEA Subscale, $F(3, 243) = 16.39, p < .001$, and Diagnosis \times TEA Subscale, $F(9, 735) = 5.35, p < .001$. Decomposing the Diagnosis \times TEA interaction with univariate Diagnosis ANOVAs for each TEA subscale revealed that the epilepsy patients (see Table 2) did not differ from the controls in the Art (95% CI = $-8.90, 6.65, p = .78$), Sensory stimuli (95% CI = $-4.47, 11.18, p = .40$), Aesthetic judgments (95% CI = $-9.99, 11.00, p = .93$), or Odor identification subscales (95% CI = $-13.82, 21.58, p = .67$). FTD patients scored lower than epilepsy patients on the Art subscale (95% CI = $-24.68, -7.59, p < .001$) and both FTD patients and AD patients scored lower than epilepsy patients on the Odor identification subscale (FTD vs. epilepsy: 95% CI = $-54.30, -15.36, p = .001$; AD vs. epilepsy: 95% CI = $-62.57, -16.32, p = .001$).

Discussion

Summary of Findings

This study sought to determine whether patients with two kinds of dementia, FTD and AD, had deficits in recognizing emotional

properties of objects as compared to healthy age-matched controls and, in an exploratory analysis, patients with epilepsy. Using the TEA, a newly developed test of ability to recognize emotional and aesthetic qualities in art and sensory stimuli, we found that patients with FTD showed significant and specific deficits in recognizing the emotions communicated by art (i.e., music and paintings); AD patients' deficits on this subtest of the TEA only approached significance. Of note, FTD patients' deficit in ability to decode emotional messages in art was quite robust, found even after controlling for the three other subscales of the TEA (Sensory stimuli, Aesthetic judgments, Odor identification) and the ability to recognize either positive emotion, negative emotion, or self-conscious emotion in other people (measured by the EFT). Both patient groups showed deficits in the ability to recognize positive emotion in other people (on the EFT), as well as deficits in identifying a specific odor. In contrast, neither FTD nor AD patients showed deficits in evaluating the pleasantness of simple sensory stimuli in normative ways, or in making aesthetic judgments for visual images.

We considered the possibility that the specific pattern of deficits shown by FTD patients in the TEA reflected item difficulty, with patients performing worst on the items that were the most difficult (i.e., aesthetic judgments of geometric shapes had the lowest percent

Table 2
Test of Emotional Aesthetics (TEA) and Empathy Film Task (EFT) Scores for Diagnostic Groups

Task	FTD	AD	Epilepsy	Controls	Total
TEA: Mean (<i>SE</i>)					
Overall	76.4 (1.8)	80.0 (1.8)	85.4 (2.8)	84.0 (.9)	82.2 (.8)
Art	76.9 (3.2)	87.5 (3.4)	93.1 (3.4)	94.2 (1.0)	90.0 (1.1)
Sensory stimuli	83.0 (2.7)	83.3 (3.0)	88.9 (3.6)	85.5 (1.2)	85.1 (1.0)
Aesthetic judgments	71.5 (2.8)	75.0 (4.1)	73.3 (5.1)	72.8 (1.8)	72.8 (1.4)
Odor identification	59.6 (6.9)	55.0 (11.4)	94.4 (5.6)	90.6 (2.3)	81.5 (2.5)
EFT: Mean (<i>SE</i>)					
Positive emotions	69.4 (5.3)	65.9 (9.7)	n/a	94.0 (1.7)	83.3 (2.5)
Negative emotions	76.2 (4.5)	93.2 (3.5)	n/a	97.2 (1.1)	90.2 (1.8)
Self-conscious emotions	44.1 (5.8)	59.1 (9.9)	n/a	80.2 (2.7)	66.7 (3.0)

Note. Values are mean percent correct on the indicated scale or subscale. FTD = frontotemporal dementia; AD = Alzheimer's disease; n/a = not available; *SE* = standard error.

Table 3
Test of Emotional Aesthetics (TEA) for FTD Subtypes

Tea component	Behavioral variant FTD	Semantic dementia
<i>N</i>	25	21
TEA: Mean (<i>SE</i>)		
Overall	77.0 (2.8)	75.0 (2.9)
Art	79.0 (4.0)	73.8 (5.8)
Sensory stimuli	84.0 (4.0)	80.2 (4.5)
Aesthetic judgments	71.2 (4.2)	73.3 (4.4)
Odor identification	56.0 (10.1)	57.1 (11.1)

Note. FTD = frontotemporal dementia; *SE* = standard error.

correct in the control sample). However, this was not the case; neither FTD nor AD patients differed significantly from controls on these items. We also considered whether the FTD patients' deficits on the TEA might be driven by particular FTD subtypes (e.g., language problems in patients with semantic dementia). However, in an exploratory analysis we found no evidence for differential performance on the TEA between patients with behavioral variant FTD and patients with semantic dementia. Finally, to examine the possibility that deficits on the TEA found in patients with FTD and AD would also be found in other neurological disorders, we examined a group of epilepsy patients awaiting surgery to treat intractable seizures. Patients with epilepsy did not show deficits on any of the TEA subscales (they did not take the EFT). Thus, the deficits we found in FTD and AD patients did not generalize to another neurological disease that has also been found to affect emotional functioning (Hixson & Kirsch, 2009).

Emotion Recognition in Neurodegenerative Disease

The present findings support and extend the existing literature on emotion recognition in neurological patients. Extensive prior evidence indicates deficits among dementia patients in the ability to recognize nonverbal expressions of emotion in other people—a crucial skill for successfully navigating the social world (e.g., Goodkind et al., 2015; Lavenu et al., 1999). Emotion recognition is subserved by complex neural circuitry (Adolphs, 2002; Preston & De Waal, 2002) and has proved to be vulnerable to a number of neurological disorders including AD and FTD (e.g., Cadieux & Greve, 1997; Goodkind et al., 2015; Gray & Tickle-Deggen, 2010; Johnson et al., 2007; Keane et al., 2002) as well as a number of psychiatric disorders (e.g., Ashwin et al., 2006; Chan et al., 2010; Rocca, et al., 2009). Neurodegenerative diseases like FTD and AD produce different patterns of deficits depending on the extent and location of neurodegeneration in large-scale brain networks (Seeley, Crawford, Zhou, Miller, & Greicius, 2009). For this reason, it is important to assess multiple aspects of emotion recognition (e.g., in objects and in people) in ways that are sensitive to patterns of lost and preserved functioning in different disorders and in different individuals.

Our findings reveal that patients with FTD and AD both show deficits in an understudied aspect of emotion recognition—the ability to decode emotional tone in works of art—but that these deficits are more profound and pervasive in FTD. These findings are consistent with prior studies that examined recognition of affective valence in art and abstract objects in FTD patients (Boutoleau-Bretonnière et al., 2016; Cohen et al., 2016; Dellach-

erie et al., 2008; Omar et al., 2011), and extend them by localizing the deficit to reading emotional tone in art, rather than appreciating the inherent pleasantness of stimuli such as scents and visual images. These deficits are particularly interesting given prior evidence that some neurological patients develop new artistic interests and skills during the course of the disease (Seeley et al., 2008).

The ability to recognize emotional tone in art is analogous to the ability to read and respond to emotions in other people, and our findings suggest that patients with FTD show distinct deficits in both. As with empathy for other humans, describing one painting or piece of music as “happier” than another requires the ability to attribute a psychological state to something other than the self (i.e., mentalizing or theory of mind). With art, this judgment may be based on a visual or auditory metaphor and may involve having a visceral reaction to the target stimulus, using that reaction to estimate the target's emotional “state.” These processes are often compromised in FTD. Prior research documents robust deficits in theory of mind, as well as deficits in comprehension of metaphor-laden, sarcastic, and nonliteral speech, in behavioral variant FTD in particular (e.g., Bora, Walterfang, & Velakoulis, 2015). Moreover, damage to structures such as the anterior insula that are critical for processing visceral feedback commonly accompanying emotion (Craig, 2009) can reduce the potency of experiential aspects of emotional responding (Verstaen et al., 2016). We and others have shown that FTD patients have problems generating robust emotional responses (Eckart, Sturm, Miller, & Levenson, 2012; Sturm, Rosen, Allison, Miller, & Levenson, 2006; Sturm, Ascher, Miller, & Levenson, 2008); a deficit that may carry over into responding to emotional messages in works of art.

In contrast, we expect that judgments about stimulus pleasantness reflect a mix of hard-wired and deeply learned associations residing in brain areas that are not early targets of FTD and AD (e.g., opioid receptor-rich regions of the nucleus accumbens and ventral pallidum; Berridge & Kringelbach, 2013). Although assessed only with a single item, our finding that FTD and AD patients had deficits in odor identification is consistent with previous findings of anosmia and other olfactory deficits in these and other neurodegenerative diseases (Luzzi et al., 2007; McLaughlin & Westervelt, 2008; Meshulam, Moberg, Mahr, & Doty, 1998). Epilepsy has been associated with impairments in aspects of emotional functioning (Hixson & Kirsch, 2009; Kanner, 2009), but in our study these did not extend to problems recognizing emotion in objects. Although epilepsy and FTD may have some overlap in regions of neuroanatomical impairment, epilepsy is more chronic than degenerative; thus, preserved neural plasticity in individuals with epilepsy may allow other regions to carry out these emotional functions.

Strengths and Limitations

The present study had several strengths, including (a) assessing emotion recognition in a number of different kinds of objects, (b) assessing multiple patient groups, and (c) exploring the robustness of found deficits (by controlling for deficits in other aspects of emotion recognition in objects and people). Limitations included (a) small sample size (especially for patients with AD who completed both the TEA and the EFT, for patients with epilepsy, and for FTD subtypes), (b) lack of power for examining ethnic differences (the sample was predominately Caucasian), (c) focusing on positive emotional judgments (i.e., happiness and pleasantness)

and not including negative emotional judgments (e.g., unhappiness and unpleasantness); and (d) differences in type of emotional judgments made for music and paintings (i.e., happiness) versus sensory stimuli and aesthetic judgments (i.e., pleasantness).

Concerning emotional judgments, we would expect similar findings if participants were asked to recognize negative emotional qualities rather than the positive emotional qualities we assessed. However, this clearly needs to be tested empirically. Similarly, additional research is needed to confirm that the patterns of deficits observed in this study are found specifically when assessing ability to detect the emotion conveyed by art (e.g., happiness in music) rather than its pleasantness. This may be particularly interesting in cases where an artistic stimulus conveys a negative emotion, yet is still pleasant (e.g., some very sad music is quite lovely), or vice versa.

Conclusion

The ability to recognize emotion in people and objects plays a vital role in our lives. The current study revealed that patients with FTD, and to a lesser extent those with AD, had impairments in recognizing emotions expressed in paintings and music, as compared to patients with epilepsy and controls. Deficits in the ability to recognize emotion in art may at first seem to be a fairly rarified loss. However, it is important to realize the important role that music and the visual arts play in the arsenal of emotion regulation. We often use music to alter our emotional state in desired ways, raising our spirits and deepening the experience of life's losses. Art is often used strategically in our home and work environments to foster particular moods. Losing access to this aspect of emotion recognition can contribute to the impoverishment of emotion that is so often seen in patients with FTD, and can add to the burden and strain experienced by their caregivers (e.g., Brown et al., 2018). These findings shed additional light on the ways that aspects of emotion recognition are preserved and diminished in different neurological diseases and add greater understanding to the profound impact that FTD can have on a wide range of emotional functions.

References

- Adolphs, R. (2002). Neural systems for recognizing emotion. *Current Opinion in Neurobiology*, *12*, 169–177. [http://dx.doi.org/10.1016/S0959-4388\(02\)00301-X](http://dx.doi.org/10.1016/S0959-4388(02)00301-X)
- Adolphs, R., Tranel, D., Hamann, S., Young, A. W., Calder, A. J., Phelps, E. A., . . . Damasio, A. R. (1999). Recognition of facial emotion in nine individuals with bilateral amygdala damage. *Neuropsychologia*, *37*, 1111–1117. [http://dx.doi.org/10.1016/S0028-3932\(99\)00039-1](http://dx.doi.org/10.1016/S0028-3932(99)00039-1)
- Ashwin, C., Chapman, E., Colle, L., & Baron-Cohen, S. (2006). Impaired recognition of negative basic emotions in autism: A test of the amygdala theory. *Social Neuroscience*, *1*, 349–363. <http://dx.doi.org/10.1080/17470910601040772>
- Berridge, K. C., & Kringelbach, M. L. (2013). Neuroscience of affect: Brain mechanisms of pleasure and displeasure. *Current Opinion in Neurobiology*, *23*, 294–303. <http://dx.doi.org/10.1016/j.conb.2013.01.017>
- Bohrt, I. C., Altmann, U., & Jacobs, A. M. (2012). Looking at the brains behind figurative language—A quantitative meta-analysis of neuroimaging studies on metaphor, idiom, and irony processing. *Neuropsychologia*, *50*, 2669–2683. <http://dx.doi.org/10.1016/j.neuropsychologia.2012.07.021>
- Bora, E., Walterfang, M., & Velakoulis, D. (2015). Theory of mind in behavioural-variant frontotemporal dementia and Alzheimer's disease: A meta-analysis. *Journal of Neurology, Neurosurgery and Psychiatry*, *86*, 714–719. <http://dx.doi.org/10.1136/jnnp-2014-309445>
- Boutoleau-Brettonnière, C., Brettonnière, C., Evrard, C., Rocher, L., Mazzietti, A., Koenig, O., . . . Thomas-Antérion, C. (2016). Ugly aesthetic perception associated with emotional changes in experience of art by behavioural variant of frontotemporal dementia patients. *Neuropsychologia*, *89*, 96–104. <http://dx.doi.org/10.1016/j.neuropsychologia.2016.06.001>
- Bowers, D., Blonder, L. X., & Heilman, K. M. (1991). *Florida Affect Battery*. Tallahassee, FL: Center for Neuropsychological Studies, Cognitive Neuroscience Laboratory, University of Florida.
- Brown, C. L., Lwi, S. J., Goodkind, M. S., Rankin, K. P., Merrilees, J., Miller, B. L., & Levenson, R. W. (2018). Empathic accuracy deficits in patients with neurodegenerative disease: Association with caregiver depression. *The American Journal of Geriatric Psychiatry*, *26*, 484–493. <http://dx.doi.org/10.1016/j.jagp.2017.10.012>
- Bucks, R. S., & Radford, S. A. (2004). Emotion processing in Alzheimer's disease. *Aging & Mental Health*, *8*, 222–232. <http://dx.doi.org/10.1080/13607860410001669750>
- Cadieux, N. L., & Greve, K. W. (1997). Emotion processing in Alzheimer's disease. *Journal of the International Neuropsychological Society*, *3*, 411–419.
- Chan, R. C. K., Li, H., Cheung, E. F. C., & Gong, Q. Y. (2010). Impaired facial emotion perception in schizophrenia: A meta-analysis. *Psychiatry Research*, *178*, 381–390. <http://dx.doi.org/10.1016/j.psychres.2009.03.035>
- Chatterjee, A. (2011). Neuroaesthetics: A coming of age story. *Journal of Cognitive Neuroscience*, *23*, 53–62. <http://dx.doi.org/10.1162/jocn.2010.21457>
- Citron, F. M. M., & Goldberg, A. E. (2014). Metaphorical sentences are more emotionally engaging than their literal counterparts. *Journal of Cognitive Neuroscience*, *26*, 2585–2595. http://dx.doi.org/10.1162/jocn_a_00654
- Cohen, M. H., Carton, A. M., Hardy, C. J., Golden, H. L., Clark, C. N., Fletcher, P. D., . . . Warren, J. D. (2016). Processing emotion from abstract art in frontotemporal lobar degeneration. *Neuropsychologia*, *81*, 245–254. <http://dx.doi.org/10.1016/j.neuropsychologia.2015.12.031>
- Collier, G. L. (1996). Affective synesthesia: Extracting emotion space from simple perceptual stimuli. *Motivation and Emotion*, *20*, 1–32. <http://dx.doi.org/10.1007/BF02251005>
- Craig, A. D. (2009). How do you feel—Now? The anterior insula and human awareness. *Nature Reviews Neuroscience*, *10*, 59–70. <http://dx.doi.org/10.1038/nrn2555>
- Dellacherie, D., Ehrle, N., & Samson, S. (2008). Is the neutral condition relevant to study musical emotion in patients? *Music Perception*, *25*, 285–294. <http://dx.doi.org/10.1525/mp.2008.25.4.285>
- Eckart, J. A., Sturm, V. E., Miller, B. L., & Levenson, R. W. (2012). Diminished disgust reactivity in behavioral variant frontotemporal dementia. *Neuropsychologia*, *50*, 786–790. <http://dx.doi.org/10.1016/j.neuropsychologia.2012.01.012>
- Essick, G. K., McGlone, F., Dancer, C., Fabricant, D., Ragin, Y., Phillips, N., . . . Guest, S. (2010). Quantitative assessment of pleasant touch. *Neuroscience and Biobehavioral Reviews*, *34*, 192–203. <http://dx.doi.org/10.1016/j.neubiorev.2009.02.003>
- Fan, Y., Duncan, N. W., de Greck, M., & Northoff, G. (2011). Is there a core neural network in empathy? An fMRI based quantitative meta-analysis. *Neuroscience and Biobehavioral Reviews*, *35*, 903–911. <http://dx.doi.org/10.1016/j.neubiorev.2010.10.009>
- Fernandez-Duque, D., & Black, S. E. (2005). Impaired recognition of negative facial emotions in patients with frontotemporal dementia. *Neuropsychologia*, *43*, 1673–1687. <http://dx.doi.org/10.1016/j.neuropsychologia.2005.01.005>
- FitzGerald, T. H. B., Seymour, B., & Dolan, R. J. (2009). The role of human orbitofrontal cortex in value comparison for incommensurable objects. *The Journal of Neuroscience*, *29*, 8388–8395. <http://dx.doi.org/10.1523/JNEUROSCI.0717-09.2009>

- Folkman, S. (1997). Positive psychological states and coping with severe stress. *Social Science & Medicine*, *45*, 1207–1221. [http://dx.doi.org/10.1016/S0277-9536\(97\)00040-3](http://dx.doi.org/10.1016/S0277-9536(97)00040-3)
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-Mental State": A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, *12*, 189–198. [http://dx.doi.org/10.1016/0022-3956\(75\)90026-6](http://dx.doi.org/10.1016/0022-3956(75)90026-6)
- Friedenberg, J., & Bertamini, M. (2015). Aesthetic preference for polygon shape. *Empirical Studies of the Arts*, *33*, 144–160. <http://dx.doi.org/10.1177/0276237415594708>
- Gabrielsson, A., & Juslin, P. N. (2003). *Emotional expression in music*. New York, NY: Oxford University Press.
- Gang, D. R. (2005). Evolution of flavors and scents. *Annual Review of Plant Biology*, *56*, 301–325. <http://dx.doi.org/10.1146/annurev.arplant.56.032604.144128>
- Gonzaga, G. C., Keltner, D., Londahl, E. A., & Smith, M. D. (2001). Love and the commitment problem in romantic relations and friendship. *Journal of Personality and Social Psychology*, *81*, 247–262. <http://dx.doi.org/10.1037/0022-3514.81.2.247>
- Goodkind, M. S., Gyurak, A., McCarthy, M., Miller, B. L., & Levenson, R. W. (2010). Emotion regulation deficits in frontotemporal lobar degeneration and Alzheimer's disease. *Psychology and Aging*, *25*, 30–37. <http://dx.doi.org/10.1037/a0018519>
- Goodkind, M. S., Sturm, V. E., Ascher, E. A., Shdo, S. M., Miller, B. L., Rankin, K. P., & Levenson, R. W. (2015). Emotion recognition in frontotemporal dementia and Alzheimer's disease: A new film-based assessment. *Emotion*, *15*, 416–427. <http://dx.doi.org/10.1037/a0039261>
- Gorno-Tempini, M. L., Hillis, A. E., Weintraub, S., Kertesz, A., Mendez, M., Cappa, S. F., . . . Grossman, M. (2011). Classification of primary progressive aphasia and its variants. *Neurology*, *76*, 1006–1014. <http://dx.doi.org/10.1212/WNL.0b013e31821103e6>
- Götz, K. O., Borisy, A. R., Lynn, R., & Eysenck, H. J. (1979). A new visual aesthetic sensitivity test: I. Construction and psychometric properties. *Perceptual and Motor Skills*, *49*, 795–802. <http://dx.doi.org/10.2466/pms.1979.49.3.795>
- Gray, H. M., & Tickle-Degnen, L. (2010). A meta-analysis of performance on emotion recognition tasks in Parkinson's disease. *Neuropsychology*, *24*, 176–191. <http://dx.doi.org/10.1037/a0018104>
- Hargrave, R., Maddock, R. J., & Stone, V. (2002). Impaired recognition of facial expressions of emotion in Alzheimer's disease. *The Journal of Neuropsychiatry and Clinical Neurosciences*, *14*, 64–71. <http://dx.doi.org/10.1176/jnp.14.1.64>
- Harlow, H. F. (1958). The nature of love. *American Psychologist*, *13*, 673–685. <http://dx.doi.org/10.1037/h0047884>
- Hixson, J. D., & Kirsch, H. E. (2009). The effects of epilepsy and its treatments on affect and emotion. *Neurocase*, *15*, 206–216. <http://dx.doi.org/10.1080/13554790802632876>
- Hsieh, S., Hodges, J. R., & Piguet, O. (2013). Recognition of positive vocalizations is impaired in behavioral-variant frontotemporal dementia. *Journal of the International Neuropsychological Society*, *19*, 483–487. <http://dx.doi.org/10.1017/S1355617712001592>
- Jacobsen, T., Schubotz, R. I., Höfel, L., & Cramon, D. Y. (2006). Brain correlates of aesthetic judgment of beauty. *NeuroImage*, *29*, 276–285. <http://dx.doi.org/10.1016/j.neuroimage.2005.07.010>
- Johnson, S. A., Stout, J. C., Solomon, A. C., Langbehn, D. R., Aylward, E. H., Cruce, C. B., . . . Paulsen, J. S. (2007). Beyond disgust: Impaired recognition of negative emotions prior to diagnosis in Huntington's disease. *Brain*, *130*, 1732–1744. <http://dx.doi.org/10.1093/brain/awm107>
- Juslin, P. N., & Laukka, P. (2003). Communication of emotions in vocal expression and music performance: Different channels, same code? *Psychological Bulletin*, *129*, 770–814. <http://dx.doi.org/10.1037/0033-2909.129.5.770>
- Kanner, A. M. (2009). Psychiatric issues in epilepsy: The complex relation of mood, anxiety disorders, and epilepsy. *Epilepsy & Behavior*, *15*, 83–87. <http://dx.doi.org/10.1016/j.yebeh.2009.02.034>
- Kawabata, H., & Zeki, S. (2004). Neural correlates of beauty. *Journal of Neurophysiology*, *91*, 1699–1705. <http://dx.doi.org/10.1152/jn.00696.2003>
- Keane, J., Calder, A. J., Hodges, J. R., & Young, A. W. (2002). Face and emotion processing in frontal variant frontotemporal dementia. *Neuropsychologia*, *40*, 655–665. [http://dx.doi.org/10.1016/S0028-3932\(01\)00156-7](http://dx.doi.org/10.1016/S0028-3932(01)00156-7)
- Kessels, R. P. C., Gerritsen, L., Montagne, B., Ackl, N., Diehl, J., & Danek, A. (2007). Recognition of facial expressions of different emotional intensities in patients with frontotemporal lobar degeneration. *Behavioural Neurology*, *18*, 31–36. <http://dx.doi.org/10.1155/2007/868431>
- Lavenu, I., & Pasquier, F. (2005). Perception of emotion on faces in frontotemporal dementia and Alzheimer's disease: A longitudinal study. *Dementia and Geriatric Cognitive Disorders*, *19*, 37–41. <http://dx.doi.org/10.1159/000080969>
- Lavenu, I., Pasquier, F., Lebert, F., Petit, H., & Van der Linden, M. (1999). Perception of emotion in frontotemporal dementia and Alzheimer disease. *Alzheimer Disease and Associated Disorders*, *13*, 96–101. <http://dx.doi.org/10.1097/00002093-199904000-00007>
- Levenson, R. W. (2007). Emotion elicitation with neurological patients. In J. A. Coan & J. J. B. Allen (Eds.), *The handbook of emotion elicitation and assessment* (pp. 158–168). New York, NY: Oxford University Press.
- Levenson, R. W., & Miller, B. M. (2007). Loss of cells—Loss of self: Frontotemporal lobar degeneration and human emotion. *Current Directions in Psychological Science*, *16*, 289–294. <http://dx.doi.org/10.1111/j.1467-8721.2007.00523.x>
- Lough, S., Kipps, C. M., Treise, C., Watson, P., Blair, J. R., & Hodges, J. R. (2006). Social reasoning, emotion and empathy in frontotemporal dementia. *Neuropsychologia*, *44*, 950–958. <http://dx.doi.org/10.1016/j.neuropsychologia.2005.08.009>
- Luzzi, S., Snowden, J. S., Neary, D., Coccia, M., Provinciali, L., & Lambon Ralph, M. A. (2007). Distinct patterns of olfactory impairment in Alzheimer's disease, semantic dementia, frontotemporal dementia, and corticobasal degeneration. *Neuropsychologia*, *45*, 1823–1831. <http://dx.doi.org/10.1016/j.neuropsychologia.2006.12.008>
- McKhann, G., Drachman, D., Folstein, M., Katzman, R., Price, D., & Stadlan, E. M. (1984). Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*, *34*, 939–944. <http://dx.doi.org/10.1212/WNL.34.7.939>
- McLaughlin, N. C., & Westervelt, H. J. (2008). Odor identification deficits in frontotemporal dementia: A preliminary study. *Archives of Clinical Neuropsychology*, *23*, 119–123. <http://dx.doi.org/10.1016/j.acn.2007.07.008>
- Meletti, S., Benuzzi, F., Rubboli, G., Cantalupo, G., Stanzani Maserati, M., Nichelli, P., & Tassinari, C. A. (2003). Impaired facial emotion recognition in early-onset right mesial temporal lobe epilepsy. *Neurology*, *60*, 426–431. <http://dx.doi.org/10.1212/WNL.60.3.426>
- Meshulam, R. I., Moberg, P. J., Mahr, R. N., & Doty, R. L. (1998). Olfaction in neurodegenerative disease: A meta-analysis of olfactory functioning in Alzheimer's and Parkinson's diseases. *Archives of Neurology*, *55*, 84–90. <http://dx.doi.org/10.1001/archneur.55.1.84>
- Møller, A. P. (1997). Developmental stability and fitness: A review. *American Naturalist*, *149*, 916–932. <http://dx.doi.org/10.1086/286030>
- Morris, J. C. (1993). The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology*, *43*, 2412–2414. <http://dx.doi.org/10.1212/WNL.43.11.2412-a>
- Neary, D., Snowden, J. S., Gustafson, L., Passant, U., Stuss, D., Black, S., . . . Benson, D. F. (1998). Frontotemporal lobar degeneration: A consensus on clinical diagnostic criteria. *Neurology*, *51*, 1546–1554. <http://dx.doi.org/10.1212/WNL.51.6.1546>

- Omar, R., Henley, S. M. D., Bartlett, J. W., Hailstone, J. C., Gordon, E., Sauter, D. A., . . . Warren, J. D. (2011). The structural neuroanatomy of music emotion recognition: Evidence from frontotemporal lobar degeneration. *NeuroImage*, *56*, 1814–1821. <http://dx.doi.org/10.1016/j.neuroimage.2011.03.002>
- Ostos, M. W., Schenk, F., Baenziger, T., & von Gunten, A. (2011). An exploratory study on facial emotion recognition capacity in beginning Alzheimer's disease. *European Neurology*, *65*, 361–367. <http://dx.doi.org/10.1159/000327979>
- Preston, S. D., & de Waal, F. B. M. (2002). Empathy: Its ultimate and proximate bases. *Behavioral and Brain Sciences*, *25*, 1–20. <http://dx.doi.org/10.1017/S0140525X02000018>
- Rankin, K. P., Kramer, J. H., & Miller, B. L. (2005). Patterns of cognitive and emotional empathy in frontotemporal lobar degeneration. *Cognitive and Behavioral Neurology*, *18*, 28–36. <http://dx.doi.org/10.1097/01.wnn.0000152225.05377.ab>
- Rascovsky, K., Hodges, J. R., Kipps, C. M., Johnson, J. K., Seeley, W. W., Mendez, M. F., . . . Miller, B. M. (2007). Diagnostic criteria for the behavioral variant of frontotemporal dementia (bvFTD): Current limitations and future directions. *Alzheimer Disease and Associated Disorders*, *21*, S14–S18. <http://dx.doi.org/10.1097/WAD.0b013e31815c3445>
- Rascovsky, K., Hodges, J. R., Knopman, D., Mendez, M. F., Kramer, J. H., Neuhaus, J., . . . Miller, B. L. (2011). Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain: A Journal of Neurology*, *134*, 2456–2477. <http://dx.doi.org/10.1093/brain/awr179>
- Reynders, H. J., Broks, P., Dickson, J. M., Lee, C. E., & Turpin, G. (2005). Investigation of social and emotion information processing in temporal lobe epilepsy with ictal fear. *Epilepsy & Behavior*, *7*, 419–429. <http://dx.doi.org/10.1016/j.yebeh.2005.07.013>
- Rocca, C. C., van den Heuvel, E., Caetano, S. C., & Lafer, B. (2009). Facial emotion recognition in bipolar disorder: A critical review. *Revista Brasileira de Psiquiatria/Brazilian Journal of Psychiatry*, *31*, 171–180. <http://dx.doi.org/10.1590/S1516-44462009000200015>
- Rosen, H. J., Gorno-Tempini, M. L., Goldman, W. P., Perry, R. J., Schuff, N., Weiner, M., . . . Miller, B. L. (2002). Patterns of brain atrophy in frontotemporal dementia and semantic dementia. *Neurology*, *58*, 198–208. <http://dx.doi.org/10.1212/WNL.58.2.198>
- Rosen, H. J., Kramer, J. H., Gorno-Tempini, M. L., Schuff, N., Weiner, M., & Miller, B. L. (2002). Patterns of cerebral atrophy in primary progressive aphasia. *The American Journal of Geriatric Psychiatry*, *10*, 89–97. <http://dx.doi.org/10.1097/00019442-200201000-00011>
- Rosen, H. J., & Levenson, R. W. (2009). The emotional brain: Combining insights from patients and basic science. *Neurocase*, *15*, 173–181. <http://dx.doi.org/10.1080/13554790902796787>
- Rosen, H. J., Pace-Savitsky, K., Perry, R. J., Kramer, J. H., Miller, B. L., & Levenson, R. W. (2004). Recognition of emotion in the frontal and temporal variants of frontotemporal dementia. *Dementia and Geriatric Cognitive Disorders*, *17*, 277–281. <http://dx.doi.org/10.1159/000077154>
- Rosen, H. J., Perry, R. J., Murphy, J., Kramer, J. H., Mychack, P., Schuff, N., . . . Miller, B. L. (2002). Emotion comprehension in the temporal variant of frontotemporal dementia. *Brain*, *125*, 2286–2295. <http://dx.doi.org/10.1093/brain/awf225>
- Schwartz, D. A., Howe, C. Q., & Purves, D. (2003). The statistical structure of human speech sounds predicts musical universals. *The Journal of Neuroscience*, *23*, 7160–7168. <http://dx.doi.org/10.1523/JNEUROSCI.23-18-07160.2003>
- Seeley, W. W., Crawford, R. K., Zhou, J., Miller, B. L., & Greicius, M. D. (2009). Neurodegenerative diseases target large-scale human brain networks. *Neuron*, *62*, 42–52. <http://dx.doi.org/10.1016/j.neuron.2009.03.024>
- Seeley, W. W., Matthews, B. R., Crawford, R. K., Gorno-Tempini, M. L., Foti, D., Mackenzie, I. R., & Miller, B. L. (2008). Unravelling Boléro: Progressive aphasia, transmodal creativity and the right posterior neocortex. *Brain*, *131*, 39–49. <http://dx.doi.org/10.1093/brain/awm270>
- Shepherd, G. M. (2006). Smell images and the flavour system in the human brain. *Nature*, *444*, 316–321. <http://dx.doi.org/10.1038/nature05405>
- Shiota, M. N. (2006). Silver linings and candles in the dark: Differences among positive coping strategies in predicting subjective well-being. *Emotion*, *6*, 335–339. <http://dx.doi.org/10.1037/1528-3542.6.2.335>
- Shiota, M. N., Campos, B., Oveis, C., Hertenstein, M. J., Simon-Thomas, E., & Keltner, D. (2017). Beyond happiness: Building a science of discrete positive emotions. *American Psychologist*, *72*, 617–643. <http://dx.doi.org/10.1037/a0040456>
- Snowden, J. S., Bathgate, D., Varma, A., Blackshaw, A., Gibbons, Z. C., & Neary, D. (2001). Distinct behavioural profiles in frontotemporal dementia and semantic dementia. *Journal of Neurology, Neurosurgery and Psychiatry*, *70*, 323–332. <http://dx.doi.org/10.1136/jnnp.70.3.323>
- Stalnaker, T. A., Cooch, N. K., & Schoenbaum, G. (2015). What the orbitofrontal cortex does not do. *Nature Neuroscience*, *18*, 620–627. <http://dx.doi.org/10.1038/nn.3982>
- Sturm, V. E., Ascher, E. A., Miller, B. L., & Levenson, R. W. (2008). Diminished self-conscious emotional responding in frontotemporal lobar degeneration patients. *Emotion*, *8*, 861–869. <http://dx.doi.org/10.1037/a0013765>
- Sturm, V. E., Rosen, H. J., Allison, S., Miller, B. L., & Levenson, R. W. (2006). Self-conscious emotion deficits in frontotemporal lobar degeneration. *Brain*, *129*, 2508–2516. <http://dx.doi.org/10.1093/brain/awl145>
- Vartanian, O., & Goel, V. (2004). Neuroanatomical correlates of aesthetic preference for paintings. *NeuroReport*, *15*, 893–897. <http://dx.doi.org/10.1097/00001756-200404090-00032>
- Verstaen, A., Eckart, J. A., Muhtadie, L., Otero, M. C., Sturm, V. E., Haase, C. M., . . . Levenson, R. W. (2016). Insular atrophy and diminished disgust reactivity. *Emotion*, *16*, 903–912. <http://dx.doi.org/10.1037/emo0000195>
- Weiss, E. M., Kohler, C. G., Vonbank, J., Stadelmann, E., Kemmler, G., Hinterhuber, H., & Marksteiner, J. (2008). Impairment in emotion recognition abilities in patients with mild cognitive impairment, early and moderate Alzheimer disease compared with healthy comparison subjects. *The American Journal of Geriatric Psychiatry*, *16*, 974–980. <http://dx.doi.org/10.1097/JGP.0b013e318186bd53>
- Werner, K. H., Roberts, N. A., Rosen, H. J., Dean, D. L., Kramer, J. H., Weiner, M. W., . . . Levenson, R. W. (2007). Emotional reactivity and emotion recognition in frontotemporal lobar degeneration. *Neurology*, *69*, 148–155. <http://dx.doi.org/10.1212/01.wnl.0000265589.32060.d3>
- Zaki, J., & Ochsner, K. N. (2012). The neuroscience of empathy: Progress, pitfalls and promise. *Nature Neuroscience*, *15*, 675–680. <http://dx.doi.org/10.1038/nn.3085>
- Zhang, Y., Schuff, N., Du, A. T., Rosen, H. J., Kramer, J. H., Gorno-Tempini, M. L., . . . Weiner, M. W. (2009). White matter damage in frontotemporal dementia and Alzheimer's disease measured by diffusion MRI. *Brain*, *132*, 2579–2592. <http://dx.doi.org/10.1093/brain/awp071>

Received October 25, 2018

Revision received June 6, 2019

Accepted June 9, 2019 ■